

Library and Knowledge Services

Mental Health Current Articles Bulletin – February 2024

How to Obtain an Article

If you would like to request any articles included in this bulletin, or to be put onto the bulletin's distribution list, please contact Library and Knowledge Services at Grantham Hospital:

Tel: 01476 464321

email: library.grantham@ulh.nhs.uk

If you require an in-depth search on a topic featured in this bulletin then please contact <u>clinical.librarian@ulh.nhs.uk</u>

Abuse/Violence/Bullying

"Older people are weak": perceptions and meanings of ageing and abuse against older people

Dias I et al

Frontiers in Sociology 2024 older people

'Yes doesn't always mean yes, but no means no': exploring the perceived ambiguities in university students' experiences of sexual consent

Savoise L et al

Canadian Journal of Human Sexuality 32 (3) 270-384

How are survivors of intimate partner violence and sexual violence portrayed on social media?

Taccini F et al

Journal of Media Psychology online first Jan 24 how are survivor

Feasibility of a reconfigured domestic violence and abuse training and support intervention responding to affected women, men, children and young people through primary care

Szilassy E et al

BMC Primary Care 15 (1) 1-18

How to prepare for conversations with children about suspicions of sexual abusr? Evaluation of an interactive virtual reality training for student teachers

Krause N et al

Child Abuse and Neglect 149 (Mar) how to prepare

Disproportionate burden of violence: explaining racial and ethnic disparities in potential years of life lost among homicide victims, suicide descendents and homicide-suicide perpetrators

Gregory M et al PLoS One 19 (2) 1-28

Autism

Co-designing behavioural activation for depression for autistic adolescents: a case seriesMohamed Z et al

CCPP 2024 epub co-designing behavioural activation

Neuroaffirming services for autistic people

Chellappa S

Lancet Psychiatry 11 (2) 96-97

Insomnia in children with Autism Spectrum Disorder: A cross-sectional study on clinical correlates and parental stress

Bianca B et al

Journal of Autism and Developmental Disorders 54 (1) 46-55

Autism Spectrum Disorder and anxiety among transgender youth: use of the Social Communication Questionnaire (SCQ)

Clyde A et al

Journal of Autism and Developmental Disorders 54 (2) 615-627

Bereavement and Grief

What bereaved children want to know about death and grief

Joy C et al

Journal of Child and Family studies 33 (2024) 327-337

Pre-death grief: support for families facing bereavement

Allen D

Nursing Standard 39 (2) 67-70

Developing a best-practice agenda for music therapy research to support informal carers of terminally ill patients pre- and post-death bereavement: a world café approach

McConnell T et al

BMC Pallliative Care 23 (1) 33-48

In-hospital end-of-left care: an appreciative analysis of bereaved family feedback

Walker W et al BJN 33 (3) 126-131

Trajectories of insomnia following bereavement

De Lang T et al Sleep Medicine 114 (Feb) 159-166

Child and Adolescent Issues

Relationship between childhood trauma and nonsuicidal self-injury among adolescents eith depressive disorder: mediated by negative life events and coping style

Xie X et al

Neuropsychiatric Disease and Treatment 19 (2023) 2271-2281

Ensuring continuity of care for young people transitioning to adult mental health services: issues faced and promising approaches

Appleton R

British Journal of Mental Health Nursing 13 (1) continuity of care

Attention deficit/hyperactivity disorder (ADHD) in children: more focus on care and support, less on diagnosis

ADHD diagnoses continue to increase but Luise Kazda and colleagues argue that the push for diagnosis could be hampering access to care and support for children

BMJ 384 (8416) attention deficit

We need more real-world trials like this one (Comment on article below)

Cartwright-Hatton S, Dunn A Lancet Psychiatry 11 (3) 161-162

Digitally augmented, parent-led CBT versus treatment as usual for child anxiety problems in child mental health services in England and Northern Ireland: a pragmatic, non-inferiority, clinical effectiveness and cost-effectiveness randomised controlled trial

Cresswell C et al

Lancet Psychiatry 11 (3) 193-209

Dementia and Alzheimer's Disease

The power of a cup of tea: psychosocial interventions in dementia

Rose T, Spencer-Davies C BJCN 29 (2) 68-75

Cut-off scores for mild and moderate dementia on the Addenbrooke's Cognitive Examination-III and the Mini-Addenbrooke's Cognitive Examination compared with the Mini-Mental State Examination

McCarthy et al

BJPsych Bulletin 48 (1) 12-18

Semantic dementia: a complex and culturally influenced presentation

Cole R et al

BJPsych Bulletin 48 (1) 44-50

Improving end-of-life care for people with dementia: a mixed-methods study

Triandafilidis Z et al

BMC Palliative Care 23 (1) 1-12

Depression

The development of psilocybin therapy for treatment-resistant depression: an update

Borissova S, Rucker J

BJPsych Bulletin 48 (1) 38-44

Anxiety and depression from childhood to young adulthood: trajectories and risk factors

LoParo D et al

Child Psychiatry and Human Development 55 (1) 127-136

An empirical comparison of statistical methods for multiple cut-off diagnostic test accuracy meta-analysis of the Edinburge postnatal scale (EPDS) depression screening tool using published results vs individual participant data

Negeri Z et al

BMC Medical Research Methodology 24 (1) 28-42

"Therapy without a therapist?": The experiences of adolescents and their parents of online behavioural activation for depression with and without therapist support

Grudin R et al

European Child & Adolescent Psychiatry 33 (1) 105-114

What treatment outcomes matter in adolescent depression? A Q-study of priority profiles among mental health practitioners in the UK and Chile

Krause K et al

European Child & Adolescent Psychiatry 33 (1) 151-166

One in five, not one in 17 youth patients deteriorate during psychotherapy for depression

McAleavey A et al

European Child & Adolescent Psychiatry 33 (1) 315-318

Exercise for the treatment of depression

Bellon J

BMJ 384 (8417) exercise

Effect of exercise for depression: systematic review and network meta-analysis of randomised controlled trials

Noetel M et al

BMJ 384 (8417) effect of exercise

Disorders

Why health anxiety needs NICE clinical guidelines

Bailey R

Lancet Psychiatry 11 (3) 164-165

Pharmacological treatments for psychotic depression: a systematic review and network meta-analysis

Vincenzo O et al

Lancet Psychiatry 11 (3) 210-220

What Is Bipolar Disorder?

Bipolar disorder is characterized by mood swings ranging from depressive lows to manic highs. Bipolar disorder affects about 8 million people in the US and an estimated 40 million individuals worldwide. Symptoms of bipolar disorder typically start between age 15 to 25 years, often with an initial episode of depression.

Voelker R

JAMA Psychiatry online first bilpolar disorder

Eating Disorders

Let the body talk: preliminary data of an Open Trial of Dance Movement Therapy for Eating Disorders

Bastoni I et al

Journal of Clinical Medicine 13 (1) 5-16

Physical health metrics are essential for establishing the effectiveness of eating disorder treatment

Ayton A

Lancet Psychiatry 11 (2) 95-96

Development and preliminary validation of a novel eating disorder screening tool for vegetarians and vegans: the V-EDS

McLean C et al

Journal of Eating Disorders 12 (1) 1-16

Treating patients with eating disorders: Guidelines from the American Psychiatric Association

Practice Guidelines

Forensic Psychiatry/Psychology

Understanding abuse, mental health and substance use: the role of victimization in a jail population

Kowalski M et al

American Journal of Criminal Justice 48 (6) 1320-1342

Prisoners with severe mental health needs are too often isolated, say monitoring boards

A lack of secure mental health beds in the community means that many prisoners in England are spending long periods in isolation, during which their health and wellbeing deteriorate, a damning report from the Independent Monitoring Boards has found.

Kmietowicz Z

BMJ 384 (8415) prisoners

Psychological framework to understand interpersonal violence by forensic patients with psychosis

Lambe S et al

BJPsych 224 (2) 47-54

Engaging forensic psychiatry patients in health-care decision making

Zhao J et al

Lancet Psychiatry 11 (3) 165-167

Intellectual/Learning Disabilities

Psychotropic medicines' prevalence, patterns and effects on cognitive and physical function in older adults with intellectual disability in Ireland: longitudinal cohort study 2009-2020

Odalovic M et al

BJPsych Open 10 (2) e39

Health support of people with intellectual disability and the crucial role of support workers Nijhof K et al

BMC Health Services Research 24 (1) health support

Clinical and cost-effectiveness of an adapted intervention for pre-schoolers with moderate to severe intellectual disabilities displaying behaviours that challenge: the EPICC-ID RCT Oddruskova T et al

Health Technology Assessment 28 (6) clinical

Mental Health & Physical Activity

Physical activity participation decreases the risk of depression in older adults: the ATHLOS Population-based cohort study

Lima R et al

Sports Medicine Open 10 (1) physical activity

Playing for more than winning: exploring sports participation, physical activity and belongingness and their relationship with patterns of adolescent substance use and mental health

Halladay J et al

Drug and Alcohol Dependence 254 (2024) 111039

Physical activity interventions to improve mental health and wellbeing in university students in the UK: a service mapping study

Malagode F et al

Mental Health and Physical Activity 26 (Mar) 100563

Mindfulness/wellbeing

The impact of gardening on well-being, mental health and quality of life: an umbrella review and meta-analysis

Pantiru I et al

Systematic Reviews 13 (2024) 1-9

Mindfulness-based programs sustainable increase mental health: the role of cognitive fusion and mindfulness practice

Lantheaume S et al

European Review of Applied Psychology, 74, Issue 2, Article 100876

Effects of combining physical activity with mindfulness on mental health and wellbeing: systematic review of complex interventions

Remskar M et al

Mental Health and Physical Activity 26 (Mar) 100575

Miscellaneous

The impact of Covid-19 3 years on: introduction to the 2-23 special issue

Maunder R et al

General Hospital Psychiatry 86 (Jan) 56-57

The old psychiatrist at table

A poem)

Kravitz R

BJPsych 224 (2) 46

The Psychological Emptiness Scale: a psychometric evaluation

Herron S et al

BJPsych Open 10 (2) e42

Social prescribing: an inadequate response to the degradation of social care in mental health

Poole R, P

BJPsych Bulletin 48 (1) 30-33

Professional

Burnout in early year medical students: experiences, drivers and the perceived value of a reflection-based intervention

Prendergast M et al

BMC Medical Education online first 2024 burnout

The sexual and gender minority (LGBTQ+) medical trainee: the journey through medical education

Danckers M et al

BMC Education 24 (2024) sexual and gender minority

Experiences and support needs of consultant psychiatrists following a patient-perpetrated homicide

Hussain Q et al

BJPsych Bulletin 48 (1) 5-11

Commissioning framework for mental health inpatient services

As well as improving experience and outcomes, this will support people to strengthen their relationships, sense of belonging and connection to local services. Creating this strategic plan and achieving its aims will require strong collaboration, particularly with local authorities and public health, and the harnessing of local information such as that within the Joint Strategic Needs Assessment. This framework summarises the issues the plan will need to address as a minimum.

NHS England Feb 24

Self-harm and Suicide

Suicide within 1 year of non-fatal overdose: risk factors and risk reduction with medications for opioid use disorder

Nowels M et al

General Hospital Psychiatry 86 (Jan) 24-32

'Am I really the priority here?': help-seeking experiences of university students who self-harmed

Tickell A et al

BJPsych Open 10 (2) e40

Associations between unintentional injuries and deliberate self-harm behaviors of children and adolescents: a school-based cross-sectional survey

Liu J et L

General Hospital Psychiatry 86 (Jan) 67-74

Suicide and safety in mental health annual report: UK patient and general population data 2011-2021

This report is based on people aged 10 and above who died by suicide between 2011 and 2021 across all of the UK. In addition to findings in relation to this general population data, it also contains key messages relating to: mental health patient suicides and acute mental health care; suicide by patients following one-off assessment; suicide by autistic people and/or patients with ADHD; suicide by in-patients aged under 25; suicide by students in England and Wales; and patients who died by suicide in public locations. It also shares findings on social and clinical characteristics National Confidential Inquiry into Suicide and Safety in Mental Health (NCISH)

Therapies/OT in mental health

Forest therapy using virtual reality in the older population: a systematic review

Clemente D et al

Frontiers in Psychology epub forest therapy

Treatment with psychedelics is psychotherapy: beyond reductionism

Grunder G et al

Lance Psychiatry 11 (3) 231-236

Acupuncture for combat-related posttraumatic stress disorder\; a randomized clinical trial Hollifield M et al

JAMA Psychiatry online first acupuncture

Veterans and Mental Health

Associations between predictors of PTSD and Psychosocial Functioning in veterans: results from a longitudinal assessment study

Pearson R et al

Depression and Anxiety 2024 ID 9719536

Sleep and pain in veterans with chronic pain: effects of psychological pain treatment and temporal associations

Wilson M et al

Psychological Trauma online first Jan 24 moral injury

A controlled trial of *adaptive disclosure-enhanced* to improve functioning and treat posttraumatic stress disorder

This is a randomized controlled trial (NCT03056157) of an enhanced adaptive disclosure (AD) psychotherapy compared to present-centered therapy (PCT; each 12 sessions) in 174 veterans with posttraumatic stress disorder (PTSD) related to traumatic loss (TL) and moral injury (MI). AD employs different strategies for different trauma types. AD-Enhanced (AD-E) uses letter writing (e.g., to the deceased), loving-kindness meditation, and bolstered homework to facilitate improved functioning to repair TL and MI-related trauma

Brett Le et al

Journal of Consulting and Clinical Psychology 92 (3) 150-164

Effectiveness of the massed delivery of unified protocol for emotional disorders within an intensive outpatient program for military service members and veterans

Sherrill A et al

Psychological Services (Jan 2024) effectiveness

The effect of companion dog adoption on loneliness in veterans with posttraumatic stress disorder: an exploratory study

Adler R et al

Traumatology (Feb 2024) companion dog adoption

Bulletin prepared by LPFT Library and Knowledge Services **Tel**: 01476 464321

email: <u>library.grantham@ulh.nhs.uk</u> <u>www.nhslincslibrary.uk</u>